

EDICT®

DISEASE CONTROL

HOW TO USE

RESTRAINTS

DO NOT mow lawns for 7 days after application.

DO NOT spray when rain is expected within 12 hours.

DO NOT apply when soil is dry and plants are suffering from moisture stress.

DO NOT spray when shade temperatures are likely to exceed 30 °C.

DO NOT use on edible plants.

Shake well before measuring amount for the rate.

SITUATION	DISEASE	RATE	CRITICAL COMMENTS
Azaleas	Azalea Petal Blight	11 drops/L water or 7.5 mL/10 L	Apply thoroughly every two weeks from the expanded bud stage (before buds colour) until the end of flowering.
Ornamentals	Botrytis Blight or Grey Mould, Downy Mildew, Leaf Spots, Rusts		Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Apply as a preventative application 7-14 days apart when conditions favour disease development (e.g. warm, rainy weather), or as a curative treatment every 7-14 days after the onset of disease.
Roses	Black Spot		
Ornamentals; Roses	Powdery Mildew	22 drops/L water or 15 mL/10 L water	
Ornamentals and non-fruit- bearing plants of the Myrtaceae family (eucalypts, melaleuca, calistemon)	Myrtle Rust	11 drops/L water or 7.5 mL/10 L	Apply at first sign of disease or when conditions favour disease development. Curative: For existing infections, apply 2 applications at a 14-day interval. Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves. Preventive: Spray to run-off ensuring thorough coverage of all foliage including the underside of leaves every 14 days according to infection pressure.
Lawns	Anthracnose, Brown Patch, Winter Fusarium, Leptosphaerulina, Curvularia	Spray: 29 drops in 1 L water over 10 m ² or one entire 15 mL tube in 7.5 L water over 75 m ² . Watering can + sprinkle bar: 29 drops in 5 L water over 10 m ² .	For preventative control, apply every 21-28 days when conditions favour disease development (e.g. warm, rainy weather). For curative control, apply when disease symptoms first appear and repeat every 14-21 days. For Brown patch control use the watering can method to ensure product is washed into the crown and root zone.
	Dollar Spot	Spray: 43 drops in 1 L water over 10 m ² or one entire 15 mL tube in 5 L water over 50 m ² . Watering can + sprinkle bar: 43 drops in 5 L water over 10 m ² .	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.

An effective combination of two fungicides to prevent and control diseases on Azaleas, roses, ornamentals and lawns. Tebuconazole works systemically from within the plant to control disease and trifloxystrobin helps prevent fungal infection.

CAUTION

DO NOT ENTER treated areas until spray has dried.

DO NOT allow containers or spray to get into drains, sewers, ponds or streams.

DO NOT feed grass clippings from treated areas to poultry or other animals.

PLANT SAFETY

EDICT® has been tested on a range of different ornamental varieties and plant safety has been demonstrated. Since the susceptibility of ornamental varieties can vary, it is recommended to test the product on a small number of plants first in order to assess the level of plant safety. If lawns are nutrient deficient, slight yellowing may occur, which can be corrected by applying lawn fertiliser.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool dry place out of reach of children and pets. **DO NOT** store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

Australian Agribusiness (Holdings) Pty Ltd
ABN: 61 135 355 958
Suite 201, 3 Rider Boulevard, Rhodes NSW 2138
PH: 1800 631 008 www.aus-ag.com.au

APVMA Approval No: 81858 / 141932

SAFETY DIRECTIONS

Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

SDS

Additional information is listed in the safety data sheet (SDS). A safety data sheet for EDICT® Disease Control is available from Australian Agribusiness (Holdings) Pty Ltd on request.

CONDITIONS OF SALE:

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. No warranty (other than non-excludable statutory warranties) of merchantability or fitness for a particular purpose, express or implied, extends to the use of product contrary to label instructions, or under off-label permits not endorsed by Australian Agribusiness (Holdings) Pty Ltd or under abnormal conditions. Australian Agribusiness (Holdings) Pty Ltd accepts no liability for any loss or damage arising from incorrect storage, handling or use.

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

EDICT®

DISEASE CONTROL

ACTIVE CONSTITUENTS 200 g/L TEBUCONAZOLE
100 g/L TRIFLOXYSTROBIN

For Azaleas, Roses, Ornamentals and Lawns.
Systemic disease control.
Controls azalea petal blight, myrtle rust, rust and mildew, lawn diseases as specified in the How to Use Table.

IMPORTANT: THIS LEAFLET IS PART OF THE LABEL ATTACHED TO THE CONTAINER. READ THOROUGHLY BEFORE OPENING OR USING THIS PRODUCT.

Australian Agribusiness (Holdings) Pty Ltd

ABN: 61 135 355 958

Suite 201, 3 Rider Boulevard, Rhodes NSW 2138

PH: 1800 631 008 www.aus-ag.com.au