

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SEGWAY[®]
400 SC FUNGICIDE

ACTIVE CONSTITUENT:
400 g/L CYAZOFAMID

GROUP 21 FUNGICIDE

For the control of fungal diseases in turf as
specified in the DIRECTIONS FOR USE table.

ISK BIOSCIENCES

NET CONTENTS
1 LITRE

PEEL HERE ►

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

When opening the container, preparing the product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and wear elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre.
Phone Australia 131 126; New Zealand 0800 764 766.

ISK ISK Biosciences Oceania Pty Ltd

Distributed in Australia by:

Amgrow Australia Pty Ltd

ABN 22 069 900 456

82 Christensen Rd, Stapylton QLD 4207

Customer Service Number: 1800 063 619

APVMA Approval No.: 86803/116628

Batch No.:

D.O.M.:



9 319943 009362

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

SEGWAY[®]

400 SC FUNGICIDE

ACTIVE CONSTITUENT:
400 g/L CYAZOFAMID

GROUP	21	FUNGICIDE
-------	-----------	-----------

For the control of fungal diseases in turf as specified in the DIRECTIONS FOR USE table.

ISK BIOSCIENCES

NET CONTENTS
1 LITRE

RESTRAINTS

DO NOT apply through any type of irrigation equipment.

DO NOT apply with aircraft.

SPRAY DRIFT RESTRAINTS

DO NOT apply with spray droplets smaller than a MEDIUM spray droplet size category according to nozzle manufacturer specifications that refer to the ASAE S572 Standard or the British Crop Production Council Guideline.

DO NOT apply when wind speed is less than 3 or more than 20 kilometres per hour as measured at the application site.

DO NOT apply during surface temperature inversion conditions at the application site.

Users of this product **MUST** make an accurate written record of the details of each spray application within 24 hours following application, and must **KEEP** this record for at least 2 years.

The spray application details that must be recorded are:

1. date with start and finish times of application;
2. location address and paddock(s) sprayed;
3. full name of this product;
4. amount of product used per hectare and number of hectares applied to;
5. crop or situation and weed or pest;
6. wind speed and direction during application;
7. air temperature and relative humidity during application;
8. nozzle brand, type, spray angle, nozzle capacity and spray system pressure measured during application;
9. name and address of person applying this product.

(Additional record details may be required by the state or territory where this product is used.)

MANDATORY NO-SPRAY ZONES

DO NOT apply if there are aquatic and wetland areas including aquacultural ponds, surface streams and rivers downwind from the application area and within the mandatory no-spray zones shown in the table below.

No-spray zones for protection of the aquatic environment:

Crop Downwind mandatory no-spray zone	Turf 5 metres
---------------------------------------	---------------

DIRECTIONS FOR USE

Crop	Disease	Rate	WHP	Critical Comments
Turf (golf and bowling greens, other intensively managed turf)	Pythium leaf blight Pythium root dysfunction (<i>Pythium</i> spp.)	10 mL / 100 square metres (1.0 L / Ha)	-	When conditions favour disease development, apply consecutive sprays of this product at approximately 21 day intervals. Apply as a foliar spray. DO NOT wait for disease to appear. Apply Segway 400 SC in 400 L to 500 L water per hectare.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

Turf: DO NOT GRAZE TREATED TURF OR LAWN; OR FEED TURF OR LAWN CLIPPINGS FROM ANY TREATED AREA TO POULTRY OR LIVESTOCK.

GENERAL INSTRUCTIONS

Mixing

Fill minimum 50% of the required water into the spray tank, and agitate when adding the required amount of Segway 400 SC Fungicide. Finally add the rest of the required water volume. Keep the spray solution agitated until all product is applied. Never prepare more spray solution than required.

Compatibility

For information on the compatibility of Segway 400 SC Fungicide with other products, contact your local ISK Biosciences Oceania Pty Ltd representative.

FUNGICIDE RESISTANCE WARNING

GROUP	21	FUNGICIDE
--------------	-----------	------------------

For fungicide resistance management Segway 400 SC Fungicide is a group 21 fungicide. Some naturally occurring individual fungi resistant to the product and other group 21 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by the product or other Group 21 fungicides, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, ISK Biosciences Oceania Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT graze or feed treated crops or turf to animals.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment that may cause drift onto nearby plants/crops, cropping lands or pastures.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

SAFETY DIRECTIONS

When opening the container, preparing the product for use and using the prepared spray, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and wear elbow length chemical resistant gloves. Wash hands after use. After each day's use wash gloves and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre.
Phone Australia 131 126;
New Zealand 0800 764 766.

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet that is available from the supplier.

CONDITIONS OF SALE

The use of ISK Biosciences Segway 400 SC Fungicide being beyond the control of the manufacturer, no warranty expressed or implied is given by ISK Biosciences Oceania Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and ISK Biosciences Oceania Pty Ltd accepts with no responsibility for any consequences whatsoever resulting from the use of this product.

Additional information required under the Globally Harmonised System (GHS) classification of the substance/mixture:

Very toxic to aquatic life. Very toxic to aquatic life with long lasting effects.



ISK Biosciences Oceania Pty Ltd

Distributed in Australia by:

Amgrow Australia Pty Ltd

ABN 22 069 900 456

82 Christensen Rd, Stapylton QLD 4207

Customer Service Number: 1800 063 619